

The Effects of Selective Aerobic Exercise on Cardiovascular Fitness in DebreMarkos Preparatory School Sedentary Female Students

FentaBitew*

Department of Sport Science, DebreMarkos University, DebreMarkos, Ethiopia

Email: fentabit@gmail.com

Abstract

The study was conducted to investigate the effect of aerobic exercise on improving cardiovascular fitness of DebreMarkos preparatory school at DebreMarkos town. 15 sedentary preparatory school female students were selected as the study subjects and their age range were from 16-18 years. All selected participants were participated in moderate intensity aerobic exercise for 8 consecutive weeks (3 days per week), 60 minutes duration per day. 12 minutes run test and step test were used and pre, during and after training test were conducted on cardiovascular fitness variables. The data collected from the study was analyzed quantitatively using descriptive statistics of central tendency of mean and standard deviation and determined the difference between initial and final mean for participants. According to the analyzed data in 12 minutes run test 826.86 meter mean difference was recorded and in step test 28.13 minute per bet mean difference was recorded and also in 12 minutes run test 73.33% of participants located under good performance zone and 26.66% of participants located under marginal zone after training. Based on the findings, it can be concluded that moderate aerobic exercise has positive effect on cardiovascular fitness of sedentary female students.

Key words: Aerobic exercise, improving cardiovascular fitness, sedentary females

1. INTRODUCTION

1.1. Background of the Study

Aerobic exercise is an activity that strengthens the lungs and heart and it improves your body's ability to use oxygen. It can improve your heart rate and blood pressure as well as your breathing^[1]

Cardiovascular fitness is important for all human beings irrespective of their age. According to ^[2]a given work may not be carried out if the required physical strength is not available. Fitness is the first and foremost things to enjoy the life fully. Regular physical activity; fitness and exercise are critically important for health and wellbeing of people of all whether they are

participated in vigorous exercise or some type of moderate health enhancing physical activity. Even for every old adult, mobility and functioning can be improved through physical activity^[3]

Aerobic activity should be an integral part of every exercise prescription for an apparently health individual, but how much aerobic in relation to resistance training depends in large part on each client's current condition and his or her fitness goals. Whether working with someone who was previously sedentary or a world class athlete's fitness related goals should be precise possible and future objectives. Measurement to make to insure the clients is progressing towards those goals whether

they are fat reduction, general fitness or competitive athletes ^[4]

Appropriate regular daily physical activity is a major component in preventing chronic disease, along with a healthy diet and not smoking for individuals. Daily physical activity is a powerful means of preventing chronic disease. A reliable experience and scientific evidence show that regular physical activity provides people both male and female of any conditions including disabilities with a wide range of physical, social and mental health benefits ^[5]

Aerobic exercise is some time known as cardio exercise that requires pumping of oxygenated blood by the heart to deliver oxygen to working muscle and stimulates the heart rate and breathing rate to increase in a way that can be sustained for the exercise session. There are different of aerobic exercise such as cardio machine, spinning, running, swimming, walking, jogging and dancing are aerobic exercise that can help to prevent the chance of developing some cancers, diabetes, depression and cardio vascular disease ^[6]

Regular aerobic exercise will produce beneficial effects for many age groups providing the exercise is specific and appropriate to the level of fitness of the individual. According to ^[7]progressive exercise correctly performed will increase the level of fitness and improve health. It will also create a sense of wellbeing produce greater energy and reduce the risk of developing many disease. Exercise makes demands on the body system over and above normal every day activities and as the result the system adopt anatomically and physiologically ^[8]

Depending on the above scholars and reviews idea the main brain storming and interest of the researcher to conduct this research is investigated the effect of selective aerobic exercise on cardio vascular fitness on DebreMarkos preparatory school sedentary female students.

1.2. Statement of the problem

Aerobic exercise is fundamentally important to conduct screening for health concerns especially for cardio vascular fitness, live reducing blood pressure, improving circulation efficiency, reducing the risk of diabetes and facilitating the flow of air in and out of lungs. Many

research studies say physical exercises are important for the development of all physical fitness. But no research was done in DebreMarkos preparatory school on female cardio related physical fitness problems.

Now a days in our country Ethiopia, because of sedentary life style most people are attacked by chronic disease such as diabetes, hypertension, coronary heart disease, asthma and some other upcoming disease. According to many research studies findings, physical inactivity is one of the causes for the development of chronic disease and poor health fitness. Similarly, in DebreMarkos Preparatory school students are living sedentary life style due to poor culture of having regular physical exercise.

Therefore, the research tried to investigate the effect of selective aerobic exercise on cardio vascular fitness on DebreMarkos preparatory school sedentary female students on those who continually conduct physical exercise by preparing fitness training plan in month and the result was examined through experimentation.

1.3. Basic research questions

The following research questions were raised to achieve the objective of the research.

1. Does aerobic exercise affect cardiovascular endurance?
2. What is the relationship between aerobic exercise and cardiovascular endurance?
3. How aerobic exercises affect cardiovascular endurance?

1.4. Objective of the study

1.4.1. General Objective

The overall aim of the research is to investigating the effect of aerobic exercise on cardiovascular endurance on DebreMarkos Preparatory school sedentary female students.

1.4.2. Specific objectives

The following specific objectives were addressed in this research

1. To test the effect of aerobic exercise on cardiovascular fitness on DebreMarkos preparatory school sedentary Female students.

2. To evaluate the relationship between selective aerobic exercise and cardiovascular fitness.
3. To find out how selective aerobic exercise affects cardiovascular fitness.

1.5. Significance of the study

The research has the following significances:

1. The primary significant of the research was the researcher confidently knows the effect of aerobic exercise on cardiovascular fitness.
2. Sedentary female students in DebreMarkos Preparatory school to understand the effect of aerobic exercise on cardiovascular fitness.
3. A base line for further studies on aerobic exercise on cardiovascular fitness.

2. Methods and Materials

2.1. Study area and period

The study has been conducted in DebreMarkos Town, which is located at North West of the capital city Addis Ababa with 300km distance and 265km south east of Bahir Dar; the capital of Amhara national regional stat. the town has 1380ml average annual rainfall and minimum and maximum temperature of 15⁰C and 22⁰C respectively. The study period was took place in 2011E.C/2019GC.

2.2. Study Design

Pre-test and post-test experimental research design was used. To arrive at the final conclusion and to examine whether selective aerobic exercise have positive effects on cardiovascular fitness or not, pre exercise during exercise and post exercise experimental tests were taken by the respective employed physical quality measurements. Data was collected from primary source.

2.3. Population of the study

The targeted populations of the study were 45 female students in DebreMarkos preparatory school. Hence, the study was concerned with the examination of effects of aerobic exercise on cardiovascular endurance in sedentary female students.

2.4. Sample size and Sampling techniques

Among the targeted population of 45 students total number of females who willingly registered to participate in this study was 15. Based on medical history, questionnaires as well as inclusion and exclusion criteria 30 participants were excluded and 15 sedentary female students were fulfill inclusion criteria. Inclusion and exclusion criteria in a sense those participants who were healthy according to medical history questionnaire and who were included the age of 16-18 years where included in this study because majority of students were found in this age. However, those who did not qualify were excluded. Therefore, 15 sedentary female students in DebreMarkos preparatory school were selected as subject unit. Purposive sampling technique was used.

2.5. Source of data & data collection instrument

Primary source of experimentation on result was a source of data. The collected through cardiovascular fitness test especially 12 minutes run test and step test. All the participants were instructed the first two days to run 12 minutes without training and record the result of each participant, and also the second two days the participants were instructed step up and down on 12 inch bench for three minutes at a rate of 24 steps per minute and recorded the second of the participants.

After measuring cardiovascular fitness before training by using 12 minutes run test and step test, start a training for 8 consecutive weeks by preparing training plan and measured cardiovascular endurance during training for 4 weeks of aerobic exercise using 12 minutes run test and step test and recorded the score of the participants. After giving 2 month training, measured the cardiovascular fitness by 12 minutes run test and step test and recorded the source of the participants. Finally, the result of the cardiovascular endurance before training and after training was compared.

Table I: The study design layout

No	Treatment	Aerobic Exercise program
1	Frequency	3 days per week
2	Total duration	8 weeks
3	Duration/session	40-60 minutes
4	Intensity	Moderate (55-69 HR max)
5	Exercise days	Thursday, Sunday and Tuesday
6	Time of training	Morning

2.6. Data Analysis

Descriptive statistics of mean and standard deviation were used and quantitative data analysis techniques were used for the research. The collected data were analyzed, interpreted and tabulated into meaning full idea using tables with appropriate standards, and percentage.

2.7. Ethical consideration

Initially letter of permission was taken from DebreMarkos University, Department of Sport Science and submitted to DebreMarkos preparatory school. Participants on the study were willing and the researcher was explained the purpose of the study for every participant and obtain information consent before beginning of the test. Moreover, culture and traditional value of all the population were taken to an account.

2.8. Measures

The particular employed physical quality measurements took a pre-test and post-test were took place whether selective aerobic exercise were positive effects on cardiovascular fitness or not, pre-exercise during practice and post-exercise experimental tests. In order to now the effect of aerobic exercise on cardiovascular fitness, an international standard of 12-minute and step test standards were employed.

Table II: Standards of 12-minute run test

Classification	Age Categories			
	17-26	27-39	40-49	50+
High performance zone	2320+	2160+	2000+	1840+
Good fitness zone	2000-2319	1920-2159	1840-1999	1680-1839
Marginal zone	1840-1999	1680-1919	1600-1839	1520-1679
Low zone	<1840	<1680	<1600	<1520

Adopted from Concept of physical fitness for wellness 10th edition, Charles B. Corbin et al., (2000)

Table III: Standard of the step test

Classification	60-second heart rate
The high –performance zone	84 or less
Good fitness zone	85-95
Marginal zone	96-119
Low zone	120 and above

Adopted from Concept of physical fitness for wellness 10th edition, Charles B. et al., (2000)

3. Result

To achieve the purpose of the study 15 female students from DebreMarkos Preparatory school sedentary female students were selected as a subjects and their age was 16-18 years. Selective aerobic exercise was given for 8 consecutive weeks. The variables selected for this study were cardiovascular fitness. Pre, during and post-test were conducted for all 15 subjects on cardio vascular fitness and the score were recorded and then the collected data was analyzed quantitatively by descriptive statics of central tendency. The results for cardiovascular fitness are discussed below.

Table IV: Results of 15 subjects’ in step test standards before, during and after training

Standard classification	Number of subject before training	%	Number of subjects during training	%	Number of subjects after training	%
High performance zone	-	-	-	-	-	-
Good fitness zone	-	-	-	-	11	73.33
Marginal zone	-	-	-	-	4	26.66
Lower zone	15	100	15	100	15	100

According to table iv, there was no subject who has high performance, good fitness and marginal zone level before and during training. All participants were located in low zone but after training 73.33% of the participants located under good fitness zone and 26.66% of participants located under marginal zone. This result indicated that effective change was observed on participant’s cardiovascular level when they are engaged in selective aerobic exercise.

Table V: Mean value of 12 minute run and step test of sedentary female students.

Dependent variables	Mean value before training	Mean value during training	Mean value after training
12 minute run test	1200.2	1708.46	2027.06

Step test	123.06	101.33	94.93
-----------	--------	--------	-------

According to table v: there was significant difference between pre to post-test score and performance was due to aerobic exercise in which they were engaged in. the mean score value for 12 minute run test before selective aerobic exercise was 1200.2 meter, during training test which was measured at the 4th week of aerobic exercise was 1708.46 meter and after training test which was measured after 8th week of aerobic exercise was 2027.06 meter. When we compared the mean value score before training test with the mean score value of after 8th week aerobic exercise, the mean difference value increased by 826.86 meter (69%). This result indicated that, the effective change was observed on participants' cardiovascular fitness level.

The mean score value for step test before training was 123.06min/beat, during training test was 101.33min/beat and after training test was 94.93min/beat. When we compare the mean value score before training test with the mean score value of 8 weeks aerobic exercise, the mean difference value was reduced by 28.13min/beat. This was due to 8th consecutive week's aerobic exercise under which the participants were through effective change on the participants' heart rate.

3.1. Standard Deviation

Table VI: standard deviation of the above result to mean

Dependent variables	Standard deviation before training	Standard deviation during training	Standard deviation after training
12 minute run test	310.14	441.49	523.83
Step test	31.78	26.17	24.52

According to numerical values indicated in table vi, 12 minutes run test before training mean of cardiovascular fitness is deviated positively by 310.14. during training which is measured at 4th week of aerobic exercise the mean of cardiovascular endurance is deviated positively by 441.49, and after training which was taken after a two week of aerobic exercise the mean of cardiovascular fitness is deviated positively by 523.83. From the above result we infer that selective aerobic exercise has an increasing effect on cardiovascular fitness.

In step test before training which was measured without training mean of cardiovascular fitness is deviated

positively by 31.78. During training which is measured at 4th week of aerobic exercise the mean of cardiovascular fitness is deviated positively by 26.17 and after training which was taken after 8 week of aerobic exercise the mean of cardiovascular fitness is deviated positively by 24.52. This result indicates that the effective change was observed on participants cardiovascular fitness parameters of heart rate.

4. Discussion

To achieve the purpose of the study 15 female students from DebreMarkos Preparatory school sedentary female students were selected as a subjects and their age was 16-18 years. Selective aerobic exercise was given for 8 consecutive weeks. The variables selected for this study were cardiovascular fitness. Pre, during and post-test were conducted for all 15 subjects on cardio vascular fitness and the score were recorded and then the collected data was analyzed quantitatively by descriptive statics of central tendency.

A sedentary life style is a type of life style with little or no physical activity. A person living a sedentary life style is often sitting or lying down while engaged in an activity like reading, socializing, watching television, playing video game, or using mobile phone or computer for much of the day. A sedentary life style can potentially contribute to ill health and many preventable cause of death.

^[9]State that, active people increase their life expectancy by two years compared to those who are inactive. Sedentary people experience a 20 percent to two-fold increase in early death compared to active people. They all so notes that physical inactivity, in combination with poor eating patterns, ranks with tobacco use among the leading preventable contributors to death. Sedentary life style and lack of physical activity can contribute for risk factors for anxiety, cardiovascular disease, breast cancer, depression, diabetes and high blood pressure

^[10]State that, aerobic exercise is a form of exercise that requires a continuous use of oxygen over an extended period of time. This usually means at list 15 to 30 minutes of continuous exercise and aerobic exercise helps to develop cardio vascular endurance.

^[1]Recommend that, aerobic exercise always be performed at your target heart rate for cardiovascular benefit. The target heart rate is between 60 to 90 percent of the difference between your resting heart rate and your maximum heart rate. Maximum heart rate is 220 minutes your age and an average resting heart rate is about 70. To achieve and maintain cardiovascular endurance people should exercising 3 to 5 days per week. The exercise session should include 15 to 60 minutes of continuous aerobic activity. The number of minutes depends on the intensity of the activity.

The American college of Medicine tends to favor aerobic activity of longer duration and less intensity. When this plan is followed, there is more likely to be a total fitness effect and less likely to be any health hazards ^[10]. Aerobic exercise are cardio machines, spinning, running, distance swimming, speed walking, jogging aerobic dancing bicycling and rope skipping. According to ^[11] aerobic exercise can prevent the chance of developing some cancers, diabetes, depression, cardiovascular disease and osteoporosis

Aerobic exercise has many benefits for sedentary person. ^[1]state that, strengthening muscles involved in respiration to facilitate the flow of air in and out of lungs and strengthening and enlarging heart muscle to improve its pumping efficacy and reduce resting heart rate; improving circulation efficiency and reducing blood pressure and increase the total number of red blood cells in the body, facilitating transport of oxygen, improve mental health; including reducing stress and lowering the incidences of depression and reducing the risk for diabetes; improving the ability of muscles to use fats during exercise and enhancing the speed at which muscle recover from high intensity exercise and increase storage of energy molecules such as fat and carbohydrates with in the muscles allowing for increase endurance.

Cardiovascular fitness is affected by many physiological parameters including heart rate, stroke volume, cardiac output and maximum oxygen consumption. Understanding the relationship between cardio respiratory and endurance training require a review of changes that occur with increased aerobic or an aerobic capacity. An early review by ^[12]As aerobic capacity increase general metabolism rises, muscle metabolism is

enhanced, hemoglobin raise, buffers in blood stream increase, venous return is improved and stroke volume is improve.

According to ^[8]the result of cardiovascular fitness conditioning has a direct positive effect on muscular endurance and indirect effects on strength and flexibility. To facilitate optimal delivery of oxygen, the working muscles of a person need to train or participate in activities that will build up the energy stores needed for sport

Aerobic exercise can give long term effects on your cardio respiratory system. ^[13]has mention on the effect of aerobic exercise can be an effective way to increase the endurance of your cardio respiratory system by maintaining the aerobic phase of exercise for many ways.

5. Conclusion

Based on the major finding of the study, effect of selective aerobic exercise on cardiovascular fitness:

- Moderate aerobic exercise has positive effect on improvement of cardiovascular fitness.
- Regular physical exercise depend on training plan is an increasing effect on cardiovascular fitness.
- Selective aerobic exercise (walking, jogging, rope skipping, long distance running, bicycling, aerobic dance) have significant effect bon improvement of health and wellbeing of sedentary people.
- Aerobic exercise has a positive effect on cardio vascular fitness and this positive effects leads to increase stroke volume at rest, lower resting heart rate and lower blood pressure.
- Aerobic exercise has a positive effect on reducing the risk of chronic disease such as hypertension, diabetes, asthma and other upcoming disease.

Acknowledgment

I would like to tank to God forever to his sacred name who help me to incept and compile the final research article interestingly.

REFERENCES

[1].Lippincott Williams &Wikins .Handbook American College of Sports Medicine, USA.2013.

[2].Reddy. M.Comparison of Circuit Training Methods of Performance variables, USA, **2012**.

[3]. Butler et al., Physical fitness: benefits of exercising for the older patient. *Geriatrics*, **1993**(10):46-62

[4]. Ekkekakis, P., Hall, E. E. and Petruzzello, S. J.. Practical markers of the transition from aerobic to anaerobic metabolism during exercise: rationale and a case for affect-based exercise prescription. *Prev.***2004**.Med. 38, 149-159.

[5]. WHO.Global Strategy on Diet Physical Activity &Health. Accessed on October20/**2003**.

[6]. Brick. L.Fitness Aerobic, Fitness Spectrum, and Human Kinetics. USA. **1996**.

[7] Thomas D. et.al. Feet and well (4th Ed) core concept and labs in physical fitness and Wellness Mayfield Publishing Company. USA, **2001**, p45-70

[8]. Rosser.M .Body Fitness &Exercise (2nd ed) the USA, **2001**.

[9]. Chararles B. etal. Concept of PhysicalFitness. Active life style for wellness (10thed) McGraw-Hill Higher Education.USA. **2000**, p106

[10]. Linda and Philip Health a Wellness Approach Merrill published. USA. **1991**.

[11]. Heyward, V.H . Advanced Fitness Assessment and Exercise Prescription (5thed) human kinetics. USA, **2006**.

[12]. Frank Moran. Glossary of Fitness and Health Terms. USA,**2010**.

[13]. Linsa Johnson. Effect of Aerobic Exercise Live Health. University of Alabama Birmingham, USA.**2009**.