RESEARCH ARTICLE OPEN ACCESS

Evaluating the Synergistic Impact of Planting Distance and Rhizome Weight on Yield Efficiency and Quality Parameters of Ginger (Zingiber officinale Rosc.)

[1]Dr. Shraddha Tiwari, [2]Dr. Rajshree Pandey
[1]Department of Botany, Govt. T.R.S college, Rewa (M.P.)
[2]Professor of Botany, Govt. Girls P.G. college, Rewa (M.P.)
Email-shraddhatiwari114@gmail.com

Abstract:

Ginger (Zingiber officinale Rosc.) is an economically significant spice crop widely cultivated in tropical and subtropical regions. Its yield and quality parameters are influenced by multiple agronomic factors, notably the weight of seed rhizomes and the spacing between plants. This research investigates the synergistic effects of different rhizome weights (20g, 30g, and 40g) and planting distances (20×15 cm, 25×20 cm, and 30×25 cm) on ginger's growth, yield, and essential quality characteristics. The study was conducted using a factorial randomized block design with three replications under controlled field conditions. Parameters including plant height, number of tillers, rhizome yield, dry matter content, and essential oil percentage were meticulously recorded. Results revealed that heavier rhizomes combined with wider plant spacing significantly enhanced growth and productivity, while intermediate combinations yielded the highest output per hectare. This study underscores the importance of balancing planting density and rhizome vigor to achieve optimal agricultural and economic outcomes in ginger cultivation.

Keywords: Ginger, Zingiber officinale, Rhizome Weight, Planting Distance, Yield, Essential Oil, Agronomic Practices, Dry Matter, Crop Quality.

1. Introduction

Ginger (Zingiber officinale Rosc.) is not just a spice—it's a legacy crop that holds a vital place in both global kitchens and traditional medicine. For centuries, it has been valued for its distinctive flavor and healing properties. Today, its commercial importance spans continents, with India, China, Nigeria, and Indonesia emerging as major producers. In India, particularly, ginger cultivation supports both the domestic spice economy and rural livelihoods, making it a crop of socio-economic importance.

The core value of ginger lies in its rhizome, which is loaded with essential oils and bioactive compounds such as gingerol, shogaol, and zingiberene. These compounds are widely recognized for their antioxidant, anti-inflammatory, and digestive properties, explaining ginger's prevalent use in herbal formulations, wellness products, and functional foods. As awareness of natural health solutions grows, so too does the demand for high-quality ginger, both in raw and processed forms.

Yet, growing ginger efficiently is not without its challenges. Farmers often face issues related to soil fertility, disease susceptibility, high input costs, and unpredictable yields. Among the various factors that affect ginger cultivation, two stand out for their direct influence on plant health and productivity: the size of the seed rhizome and the distance between plants. Seed rhizome weight acts as a vital determinant of early plant vigor by providing stored nutrients for initial growth, while spacing influences the level of competition among plants for light, nutrients, and water.

Research indicates that closer spacing can increase plant populations per unit area but may also lead to reduced per-plant performance due to overcrowding. On the other hand, wider spacing gives plants more room to grow but may lower the total yield per hectare if not optimized. Similarly, while heavier rhizomes can boost plant performance, they also increase input costs. Therefore, balancing these two variables is key to enhancing overall crop performance.

Another important dimension is sustainability. As agriculture adapts to growing environmental and economic pressures, optimizing input use while maintaining or improving yield has become critical. Strategies that improve input-use efficiency—like choosing the right rhizome size and planting geometry—can make ginger farming more sustainable and profitable. Moreover, recent research has shown that such agronomic adjustments can also improve the essential oil content and quality of ginger, which adds even more commercial value.

Despite the growing body of work on ginger agronomy, very few studies have explored the interaction effects of rhizome size and spacing in detail. This study was designed to fill that gap by evaluating how different combinations of rhizome weight and planting distance affect ginger's growth, yield efficiency, and quality traits like essential oil content. The goal is to offer practical, data-backed recommendations that farmers can adopt to maximize their returns. Ultimately, the findings of this research aim to contribute to smarter, more adaptive cultivation strategies that align with both farmer needs and market demands.

2. Materials and Methods

2.1 Experimental Site and Soil Conditions

The experiment was conducted at an agricultural research farm situated in a subtropical zone, characterized by average annual rainfall of 1100 mm and mean temperatures ranging from 25°C to 32°C. The soil was loamy, well-drained, and slightly acidic with a pH of 6.2.

2.2 Treatment Details and Design

A factorial randomized block design (FRBD) with three replications was adopted. The treatments comprised three rhizome weights (20g, 30g, 40g) and three planting distances (20×15 cm, 25×20 cm, and 30×25 cm), totaling nine treatment combinations. Each plot measured 3m × 2m with 0.5m spacing between plots.

2.3 Cultural Practices and Crop Management

Before planting, seed rhizomes were treated in hot water at 50°C for 10 minutes to eliminate pathogens. Organic compost was applied at 20 tons per hectare. Planting was done manually in the first week of June, followed by mulching with green leaves. Drip irrigation was maintained throughout the growing season. Pest and disease management was carried out using neem-based biopesticides.

2.4 Data Collection and Statistical Analysis

Data collection included plant height at 90 and 150 days after planting, number of tillers per plant, fresh rhizome yield (g/plant), dry matter content (%), and essential oil content (%), determined using Clevenger's hydrodistillation method. Statistical analysis was performed using ANOVA, and treatment means were compared using the Least Significant Difference (LSD) test at a 5% level of significance.

Planting distance

Rhizome weight	20 g	20 × 15 cm	30 × 25 cm
	20 g 20×15 cm	30 g 25×25 cm	30 g 30×25 cm
	30 g 20×15 cm	30 g 25×25 cm	30 g 30×25 cm
	40 g 20×15 cm	40 g 25×25 cm	40 g 30×25 cm

Figure 1. Schematic diagram showing layout of treatments with different rhizome weights and planting distances.

3. Results

3.1 Growth Performance

A significant interaction was observed between rhizome weight and planting distance across all measured parameters. The treatment combination of 40g rhizome and 30×25 cm spacing resulted in the highest plant height (98.4 cm) and maximum tiller number (12.5).

3.2 Yield Parameters

This treatment also achieved the highest rhizome yield per plant (580.2 g). Intermediate treatments like 30g rhizomes at 25×20 cm showed favorable performance, especially in terms of total yield per hectare due to higher planting density.

3.3 Quality Characteristics

The same combination recorded the highest dry matter content (23.6%) and essential oil percentage (2.48%). Closer spacing with lighter rhizomes (20g at 20×15 cm) resulted in the lowest values due to increased competition and limited nutrient reserves.

3.4 Statistical Correlations

Dry matter and essential oil content were positively correlated (r = 0.87), indicating that well-spaced, vigorous plants not only yield more biomass but also exhibit superior quality traits. Economic analysis indicated that although per-plant yield was highest in wider spacing, yield per unit area was optimized in the intermediate spacing treatments.



Figure 2. Bar graph comparing average rhizome yield (g/plant) across different rhizome weight and spacing combinations.



Figure 3. Line graph showing correlation between essential oil content (%) and treatment combinations of rhizome weight and spacing.

4. Discussion

The study emphasizes that larger rhizomes store greater quantities of carbohydrates and essential nutrients, supporting vigorous early growth and canopy development. Wider spacing was shown to reduce intra-plant competition, improving photosynthetic efficiency, root development, and nutrient uptake, all contributing to superior rhizome quality.The findings align with the hypothesis that an optimal combination of rhizome weight and spacing significantly influences ginger performance. The improved dry matter and essential oil contents observed in wider spacing with larger rhizomes suggest that plants benefit from greater nutrient availability and reduced competition. This combination fosters a healthier physiological status that enhances the accumulation of essential secondary metabolites in the rhizome.

Interestingly, the 30g × 25×20 cm configuration also showed promising results. Though its per-plant yield was slightly lower than the 40g × 30×25 cm combination, its yield per hectare was competitive due to higher plant density. This suggests that economic returns can be maximized not only by focusing on per-plant productivity but also by strategically optimizing planting configuration.The study further highlights a strong positive correlation between dry matter and essential oil content, reinforcing the relationship between physiological growth vigor and biochemical quality. For farmers and agronomists aiming at both yield and quality, adopting medium to large rhizomes in moderately spaced planting schemes could offer a viable cultivation strategy.

From an agronomic perspective, the results provide evidence that tailored planting strategies can have a marked impact on overall crop success. Regional trials under varying climatic and soil conditions are recommended to fine-tune these practices for broader applicability. Additional studies on cost-benefit analyses and post-harvest characteristics can also provide deeper insights for commercial-scale adoption.

5. Conclusion

This study concludes that both rhizome weight and planting distance significantly influence the yield, growth characteristics, and quality of ginger. The 40g rhizome weight combined with 30×25 cm spacing produced the most favorable results across all measured parameters, highlighting its potential as the most productive and quality-enhancing treatment.

However, the 30g rhizome weight with 25×20 cm spacing also performed remarkably well, especially in terms of yield per hectare. This makes it a cost-effective and practical choice for small and medium-scale farmers who aim to balance input costs with output returns.The positive relationship between essential oil content and plant vigor underscores the importance of carefully managing early plant development through seed size and spacing decisions. Incorporating these findings into best practice guidelines could enhance both the agronomic efficiency and economic sustainability of ginger cultivation.Future research should explore the interaction of these factors with organic and inorganic nutrient applications, mulching practices, and climate-resilient strategies to ensure adaptability across various agro-ecological zones. This integrated approach would support the development of a holistic cultivation model for maximizing ginger productivity and quality.

6. References

1) Ahmed, S., Rahman, M. M., & Alam, M. K. (2013). Effect of planting material and spacing on growth and yield of ginger. International Journal of Agronomy and Agricultural Research, 3(9), 25–30.
2) Anburani, A., & Manivannan, K. (2002). Effect of integrated nutrient management on growth in brinjal. South Indian Horticulture, 50(4–6), 377–386.
3) Bhatt, L., & Srivastava, B. K. (2014). Influence of spacing and seed rate on ginger productivity. Indian Journal of Horticulture, 71(1), 92–95.
4) Chattopadhyay, S. B., & Dutta, S. (2005). Ginger cultivation practices in hilly regions. Spices & Aromatic Crops Journal, 10(2), 103–107.
5) Deka, B. C., & Hazarika, T. K. (2010). Ginger: Post-harvest management and value addition. Indian Horticulture, 55(2), 16–20.
6) Farooq, M., Wahid, A., & Siddique, K. H. M. (2012). Micronutrient application through seed treatments: A review. Journal of Soil Science and Plant Nutrition, 12(1), 125–142.
7) Gomez, K. A., & Gomez, A. A. (1984). Statistical procedures for agricultural research (2nd ed.). John Wiley & Sons.
8) Haque, K. M. F., & Hossain, M. A. (2010). Effect of seed rhizome size on the growth and yield of ginger. Bangladesh Journal of Agricultural Research, 35(1), 95–100.

9) Hossain, M. D., & Rahman, M. A. (2009). Agronomic performance of ginger under different spacing. Agricultural Science Digest, 29(3), 216–219.
10) Islam, M. A., & Uddin, M. N. (2009). Effect of spacing and seed rhizome weight on the yield of ginger. Journal of Agroforestry and Environment, 3(2), 1–4.
11) Khandaker, S., & Hossain, A. (2014). Influence of seed rhizome weight on ginger yield and profitability. Bangladesh Agronomy Journal, 17(2), 33–40.
12) Kumar, M., & Sinha, S. (2006). Role of plant spacing and seed weight in turmeric production. Journal of Spices and Aromatic Crops, 15(1), 50–53.
13) Mohanty, C. R., & Panda, P. K. (2011). Effect of organic manures on ginger. International Journal of Agricultural Sciences, 7(2), 472–474.
14) Nair, K. P. P. (2013). The agronomy and economy of turmeric and ginger. Elsevier.
15) Patra, N. K., & Rao, P. S. (2002). Influence of organic farming on yield and quality of ginger. Indian Journal of Agronomy, 47(1), 118–122.
16) Rahman, M. M., & Islam, S. (2015). Integrated crop management practices in ginger for sustainable production. Journal of Agricultural Extension and Rural Development, 7(7), 214–220.
17) Ravindran, P. N., & Babu, K. N. (2005). Ginger: The genus Zingiber. CRC Press.
18) Roy, R. K., & Hore, D. K. (2010). Performance of ginger genotypes under different agro-climatic zones. Indian Journal of Horticulture, 67(3), 402–406.
19) Singh, D. B., & Ghosh, S. (2003). Effect of spacing and mulching on ginger. Indian Journal of Agricultural Research, 37(4), 284–287.
20) Zaman, M. A., & Sarkar, M. A. R. (2011). Growth and yield of ginger as influenced by seed rhizome size and spacing. Journal of Environmental Science and Natural Resources, 4(2), 147–150